
Aspects of non leptonic B_s decays

R. Ferrandes

Phys. Dept. University of Bari and INFN, via Orabona 4, I-70126 Bari, Italy
 rossella.ferrandes@ba.infn.it

Non perturbative strong interaction effects make difficult a theoretical description of non leptonic weak decays of hadrons; so it is relevant the possibility to obtain predictive parametrizations of decay amplitudes. It is noticeable that the widths of a set of two-body B_s transitions can be predicted using the symmetries of QCD and available information on B decays [1]. We consider in particular a class of decay modes induced by the transitions $b \rightarrow c\bar{u}q$ ($q = d$ or s) which are collected in Table 1. These quark transitions are described by the following effective hamiltonian, obtained by a renormalization group evolution from the electroweak scale down to $\mu = m_b$: $H = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* [c_1(\mu) Q_1 + c_2(\mu) Q_2]$, where $Q_1 = (\bar{c}_i b_i)_{V-A} (\bar{d}_j u_j)_{V-A}$ and $Q_2 = (\bar{c}_i b_j)_{V-A} (\bar{d}_j u_i)_{V-A}$.

The most popular approach to compute hadronic matrix element of four quark operators is based on naive factorization [2], consisting in factorizing them in current matrix elements determined in terms of meson decay constants and semileptonic form factors. This approach has some troubles: firstly, the Wilson coefficient $c_i(\mu)$ depend on renormalization scale μ , while form factors, decay constants and physical amplitudes are scale independent. Moreover, the amplitudes are real, so that strong phases are neglected; finally, annihilation amplitudes are predicted to be tiny. Instead, data point to sizeable strong phases and to non negligible annihilation contributions, as we shall see.

An alternative approach is a model independent analysis based on flavor symmetry and experimental data. The key observation is that the various B_s decay modes are governed, in the $SU(3)_F$ limit, by few independent amplitudes [3] that can be constrained, both in moduli and in phase differences, from corresponding B decay processes. Since $\bar{B} \rightarrow DP$ decays induced by $b \rightarrow c\bar{u}d(s)$ transitions involve a weak Hamiltonian transforming as a flavor octet, using the notation $T_\nu^{(\mu)}$ for the $\nu = (Y, I, I_3)$ component of an irreducible tensor operator of rank (μ) [4], one can write: $H_W = V_{cb} V_{ud}^* T_{0-1-1}^{(8)} + V_{cb} V_{us}^* T_{-1-\frac{1}{2}-\frac{1}{2}}^{(8)}$. When combined with the initial $\bar{B}$ mesons, which form a (3^*) -representation of $SU(3)$, this leads to (3^*) , (6) and

(15*) representations, which are also those formed by the combination of the final octet light pseudoscalar meson and triplet D meson. Therefore, using the Wigner-Eckart theorem, the decay amplitudes can be written as linear combinations of three reduced amplitudes $\langle \phi^{(\mu)} | O^{(8)} | B^{(3^*)} \rangle$, with $\mu = 3^*, 6, 15^*$. By appropriate linear combinations of these amplitudes one can obtain a correspondence with the color suppressed, color enhanced and W-exchange diagrams, C , T and E , respectively, as in Table 1. The transition in the $SU(3)$ singlet η_0 involves another amplitude D in principle not related to the previous ones. The $SU(3)$ representation for B decays is reported in Table 1.

Table 1. $SU(3)$ amplitudes for B^- , $\bar{B}^0$ and $\bar{B}_s^0$ decays to $D_{(s)}P$ (P is a light pseudoscalar meson), induced by $b \rightarrow c\bar{u}d(s)$ transitions. The predicted $\bar{B}_s^0$ branching fractions are also reported.

$B^-, \bar{B}^0$	amplitude	$\text{BR}_{exp} (10^{-4})$	$\ \bar{B}_s^0$	amplitude	$\text{BR}_{th} (10^{-4})$
$D^0 \pi^-$	$V_{ud}^* V_{cb} (C + T)$	49.8 ± 2.9	$D_s^+ \pi^-$	$V_{ud}^* V_{cb} T$	29 ± 6
$D^0 \pi^0$	$V_{ud}^* V_{cb} \frac{(C-E)}{\sqrt{2}}$	2.91 ± 0.28	$D^0 \bar{K}^0$	$V_{ud}^* V_{cb} C$	8.1 ± 1.8
$D^+ \pi^-$	$V_{ud}^* V_{cb} (T + E)$	27.6 ± 2.5	$D^0 \eta_8$	$V_{us}^* V_{cb} \frac{(2C-E)}{\sqrt{6}}$	
$D_s^+ K^-$	$V_{ud}^* V_{cb} E$	0.38 ± 0.13	$D^0 \eta_0$	$V_{us}^* V_{cb} D$	
$D^0 \eta_8$	$-V_{ud}^* V_{cb} \frac{(C+E)}{\sqrt{6}}$		$D^0 \eta$		0.21 ± 0.12
$D^0 \eta_0$	$V_{ud}^* V_{cb} D$		$D^0 \eta'$		0.10 ± 0.08
$D^0 \eta$		2.2 ± 0.5	$D^0 \pi^0$	$-V_{us}^* V_{cb} \frac{E}{\sqrt{2}}$	0.010 ± 0.003
$D^0 \eta'$		1.7 ± 0.4	$D^+ \pi^-$	$V_{us}^* V_{cb} E$	0.020 ± 0.006
$D^0 K^-$	$V_{us}^* V_{cb} (C + T)$	3.7 ± 0.6	$D_s^+ K^-$	$V_{us}^* V_{cb} (T + E)$	1.8 ± 0.3
$D^0 \bar{K}^0$	$V_{us}^* V_{cb} C$	0.50 ± 0.14			
$D^+ K^-$	$V_{us}^* V_{cb} T$	2.0 ± 0.6			

We note that $\bar{B} \rightarrow D_s K$ only fixes the modulus of E , which is not small at odds with the expectations by factorization, where W -exchange processes are suppressed by ratios of decay constants and form factors and are usually considered to be negligible. What can be done is to use all the information on $\bar{B} \rightarrow D\pi$, $D_s K$ and DK (7 experimental data) to determine T , C and E (5 parameters). A similar strategy has been recently adopted in [6]. Noticeably, the combined experimental information is enough accurate to tightly determine the ranges of variation for all these quantities. In fig. 1 we have depicted the allowed regions in the C/T and E/T planes, obtained fixing the other variables to their fitted values, with the corresponding confidence levels. It is worth noticing that the phase differences between the various amplitudes are close to be maximal; this signals sizeable deviation from naive (or generalized) factorization, provides constraints to QCD-based approaches proposed to evaluate non leptonic B decay amplitudes [7, 8, 9] and points towards

large long-distance effects in C and E [10]. We obtain $|\frac{C}{T}| = 0.53 \pm 0.10$, $|\frac{E}{T}| = 0.115 \pm 0.020$, $\delta_C - \delta_T = (76 \pm 12)^\circ$ and $\delta_E - \delta_T = (112 \pm 46)^\circ$.

With the results for the amplitudes we can determine a number of B_s decay rates, and the predictions are collected in Table 1. The uncertainties in the predicted rates are small; in particular, the W -exchange induced processes $\bar{B}_s^0 \rightarrow D^+\pi^-$, $D^0\pi^0$ are precisely estimated [11]. The predicted ratio $\frac{\Gamma(B_s \rightarrow D_s^- \pi^+)}{\Gamma(B^0 \rightarrow D^- \pi^+)} = 1.05 \pm 0.24$ can be compared to the experimental value $1.32 \pm 0.18 \pm 0.38$ recently obtained by CDF Collaboration [12].

We have performed an analogous analysis for $\bar{B} \rightarrow D_{(s)}V$, $D_{(s)}^*P$ decays, for which the same $SU(3)$ decomposition holds, with amplitudes T' , C' , E' . B decay data are collected in Table 2, including the recently observed W -exchange mode $\bar{B}^0 \rightarrow D_s^* K^-$ [14], together with the predictions for B_s decays.

$SU(3)_F$ breaking terms can modify our predictions. Those effects in general cannot be reduced to well defined and predictable patterns without new assumptions. Their parametrization would introduce additional quantities [13] that at present cannot be sensibly bounded since their effects seem to be smaller than the experimental uncertainties. It will be interesting to investigate their role when the B_s decay rates will be measured and more precise B branching fractions will be available.




Fig. 1. Ratios of $SU(3)$ amplitudes obtained from B data in Table 1. The contours correspond to the confidence level of 68% (continuous line), 90% (dashed line) and 95% (long-dashed line); the dots show the result of the fit.

Acknowledgments

I thank P. Colangelo for collaboration and F. De Fazio for useful discussions.

Table 2. Experimental branching fractions of $\bar{B} \rightarrow D_{(s)}V$, $D_{(s)}^*P$ decays and predictions for $\bar{B}_s^0$ decays.

$B^-, \bar{B}^0$	$\text{BR}_{exp} (10^{-4})$	$\parallel \bar{B}_s^0$	$\text{BR}_{th} (10^{-4})$
$D^0 \rho^-$	134 ± 18	$D_s^+ \rho^-$	72 ± 35
$D^0 \rho^0$	2.9 ± 1.1	$D^0 \bar{K}^{*0}$	9.6 ± 2.4
$D^+ \rho^-$	77 ± 13		
$D_s K^{*-}$	< 9.9		
$D^0 K^{*-}$	6.1 ± 2.3	$D^0 \rho^0$	0.28 ± 1.4
$D^0 \bar{K}^{*0}$	0.48 ± 0.12	$D^+ \rho^-$	0.6 ± 2.8
$D^+ K^{*-}$	3.7 ± 1.8	$D_s^+ K^{*-}$	4.5 ± 3.1
$D^{*0} \pi^-$	46 ± 4	$D_s^{*+} \pi^-$	32 ± 2
$D^{*0} \pi^0$	2.7 ± 0.5	$D^{*0} \bar{K}^0$	4.7 ± 2.2
$D^{*+} \pi^-$	27.6 ± 2.1		
$D_s^* K^-$	0.20 ± 0.06		
$D^{*0} K^-$	3.6 ± 1.0	$D^{*0} \pi^0$	0.0057 ± 0.0017
		$D^{*+} \pi^-$	0.0115 ± 0.0035
$D^{*+} K^-$	2.0 ± 0.5	$D_s^{*+} K^-$	1.3 ± 0.2

References

1. P. Colangelo, R. Ferrandes, Phys. Lett. B **627**, 77 (2005).
2. M. Neubert and B. Stech, Adv. Ser. Direct. High Energy Phys. **15**, 294 (1998).
3. D. Zeppenfeld, Z. Phys. C **8**, 77 (1981);
B. Grinstein and R. F. Lebed, Phys. Rev. D **53**, 6344 (1996).
4. J. J. de Swart, Rev. Mod. Phys. **35**, 916 (1963).
5. S. Eidelman et al. [Particle Data Group], Phys. Lett. B **592**, 1 (2004).
6. C. S. Kim, Sechul Oh, Chaehyun Yu, Phys. Lett. B **621**, 259 (2005).
7. M. Beneke et al, Nucl. Phys. B **591**, 313 (2000).
8. C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. Lett. **87**, 201806 (2001).
9. Y. Y. Keum et al., Phys. Rev. D **69**, 094018 (2004).
10. P. Colangelo et al., Phys. Lett. B **542**, 71 (2002); Phys. Rev. D **69**, 054023 (2004); Phys. Lett. B **597**, 291 (2004); H. Y. Cheng et al., Phys. Rev. D **71**, 014030 (2005); C. K. Chua and W. S. Hou, Phys. Rev. D **72**, 036002 (2005).
11. R. Aleksan, I. Dunietz and B. Kayser, Z. Phys. C **54**, 653 (1992).
12. A. Abulencia et al [CDF]: Phys. Rev. Lett. **96**, 191801 (2006).
13. M. Gronau et al., Phys. Rev. D **52**, 6356 (1995).
14. B. Aubert et al. [BABAR], hep-ex/0604012.